

DOI: 10.5152/eurasianjmed.2019.18342

Manuscript Type: Original Article

Title: The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms

Authors: Orçun Çiftçi¹, Kerem Can Yılmaz¹, Emir Karaçağlar¹, Mustafa Yılmaz², Bülent Özin¹, İ. Haldun Müderrisoğlu¹

Institutions: 1Başkent University Faculty of Medicine, Department of Cardiology, Ankara Hospital,

Ankara, Turkey

2 Başkent University Faculty of Medicine, Department of Cardiology, Adana Hospital, Adana,

Turkey

Correspondence to: Orçun Çiftçi

You may cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

Introduction

Atrial fibrillation (AF), is a common cause of emergency department admission, particularly when associated with a rapid ventricular response [1]. AF may create myocardial ischemia or injury in the case of underlying severe CAD, but it may also mimic cardiac ischemia or infarction through elevated cardiac biomarkers and ECG changes indicative of ischemia in the absence of underlying severe or unstable coronary artery disease (CAD) [2]. Thus, there is a need for reliable predictors of severe or unstable CAD as the cause of signs and symptoms of patients with atrial fibrillation and unstable symptoms. It has been reported that CHADS₂ and CHA₂DS₂-VASc scores, which were originally developed for risk prediction of stroke and death [3-6]. could predict severe CAD in various clinical settings [7-9]. Recently, the CHA₂DS₂-VASc-FSH score was formulated, which includes hyperlipidaemia, smoking, and family history for CAD in addition to the original CHA₂DS₂-VASc score and substitutes male gender for female gender. Hence, by incorporating the majority of risk factors for CAD, this newly developed score provides a comprehensive risk assessment for CAD [9]. The aim of this study was to determine the prevalence of severe CAD and the predictive ability of the newly developed CHA₂DS₂-VASc-FSH score for the prediction of severe CAD in comparison of conventional risk scores developed for AF as well as other demographic, clinical, electrocardiographic, and biochemical parameters in patients with AF and unstable symptoms.

Materials and Methods

This study was approved by XXXXX University Institutional Review Board (Project No:KA17/104) and supported by XXXXX University Research Fund. We retrospectively analysed the medical records of 72 patients older than 18 years who presented to the emergency department of XXXXX University Faculty of Medicine Hospital between April 2011 and January 2016. All study subjects presented with symptomatic AF and a unstable clinical symptoms suggestive of cardiac ischemia, including chest pain or one or

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özın B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

more of its equivalents, including resting dyspnea, palpitations, altered consciousness, syncope, diaphoresis, agitation, decompensated heart failure or pulmonary edema, and all of them underwent invasive coronary angiography in the index hospitalization.

AF was classified into the categories of new-onset, paroxysmal, persistent, and permanent AF categories that were determined according to the 2014 American College of Cardiology/American Heart Association/Heart Rhythm Society (AHA/ACC/HRS) task force on practice guidelines, and the 2010 European Society of Cardiology (ESC) Guidelines [10,11]. Rapid ventricular response was defined as resting heart rate of greater than 100/min during AF at rest. Severe chronic renal disease referred to severely (GFR \leq 30 ml/min) or very severely (GFR \leq 15 ml/min and/or receiving renal replacement therapy) reduced renal function (stage 4 and 5 chronic kidney disease, respectively). Previous coronary artery disease was defined as invasively or non-invasively detected coronary luminal irregularities, plaques, or insignificant narrowings (\leq 50%) previously treated medically, and also significant (\geq 50%) lesions previously treated with either medical therapy, percutaneous coronary intervention, or coronary bypass surgery. Family history for coronary artery disease was defined as having a first degree relative having CAD before the age of 65 for women and 55 for men. History of smoking included past or current smoking. History of heart failure included past or current signs and symptoms of heart failure with both low ($<$ 40%) and preserved ejection fraction (\geq 40%) and/or other objective evidence of cardiac dysfunction. Hypertension was defined as ongoing therapy for hypertension, systolic blood pressure (SBP) of \geq 140mmHg or diastolic blood pressure (DBP) of \geq 90mmHg; diabetes mellitus was defined as a fasting blood glucose level $>$ 126 mg/dL or blood glucose \geq 200 mg/dL or using antidiabetic drugs; hyperlipidemia was defined as hypercholesterolemia (serum TC $>$ 5.72 mmol/L), high levels of LDL-c ($>$ 3.1 mmol/L), low levels of HDL-c ($<$ 0.9 mmol/L), hypertriglyceridemia (serum TG $>$ 1.70 mmol/L), or taking antihyperlipidemic medications. History of ischemic cerebrovascular accident was defined as having a history of ischemic transient ischemic attack or ischemic

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

cerebral stroke. Previous non-cerebral embolic events were defined as thromboembolic events to any organ other than the brain.

Cardiac biomarkers of injury, namely mass CK-MB and troponin I (cTnI), were measured both at admission and at 3 hours of follow-up. cTnI testing was performed with Architect STAT troponin I analyzer (Abbott Diagnostics) with 99th percentile cut-off value of 0.01 µg/L. Numerical troponin I values and binary (present-absent) troponin I positivity were recorded at admission and at follow-up. Troponin positivity referred to a troponin I measurement above the upper reference limit (URL). A delta troponin I change between the admission and follow-up troponin I values was recorded for every patient and defined by either as a 50% increase in the follow-up troponin I value compared to the admission value when the latter was below the upper reference limit (URL), provided that either of the measurements was above the URL; or a 20% increase in the follow-up troponin I value when the admission value was above URL [12]. Renal function tests, electrolytes, and complete blood count parameters including leucocyte count, neutrophil count, lymphocyte count, red cell distribution width (RDW), hemoglobin level, platelet count, platelet/lymphocyte ratio (PLR), and neutrophil/lymphocyte ratio (NLR) were also measured.

Electrocardiographic changes suggestive of myocardial ischemia or injury were analyzed from 12-lead ECG recordings made at admission and 3-6 hour later, and were defined as ≥ 1.0 mm horizontal or downsloping ST segment depression 0.08 seconds after the J point, symmetrical T inversion in at least 2 contiguous precordial leads excluding V1 and V2, newly developed pathological Q waves defined as initial negative waves at least 0.04 sec in duration and 0.1 mv or $> \frac{1}{4}$ of the preceding R wave in size, or newly developed loss of R wave progression across the precordial leads. Echocardiographic examinations were performed using Vivid 7 Vingmed echocardiography device (Horten, Norway). Echocardiographic left ventricular ejection fraction (LVEF) was determined by the Modified Simpson's method and systolic wall motion abnormality for each left ventricular segment was assessed by 2-D echocardiography. Left ventricular hypertrophy

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

was defined as interventricular septal and/or posterior left ventricular wall thicknesses exceeding 11 mm.

All patients in this study underwent invasive coronary angiography at consulting physician's discretion, on the basis of signs of symptoms suggesting acute cardiac ischemia or infarction. A significant coronary lesion was defined as luminal narrowing of $\geq 50\%$ in any major epicardial vessel or one of its branches. In patients with previous PCI, significant coronary lesions were defined as significant stent restenosis ($\geq 50\%$) in previously stented vessels or de novo significant luminal narrowing ($\geq 50\%$) in previously untreated vessels. In patients who underwent a prior coronary bypass grafting operation, severe coronary artery disease was defined as significant de novo lesions in bypass grafts supplying one or more epicardial vessels, in non-bypassed coronary vessels, or in native vessels at sites distal to the insertion sites of bypass grafts. A non-significant coronary lesion was defined as luminal irregularities or plaques, or non-severe coronary lesions not exceeding 50%. The study population was classified into two groups depending on coronary angiography on invasive coronary angiography, and the two groups were compared with respect to the CHADS₂, CHA₂DS₂-VASC, and CHA₂DS₂-VASC-FSH scores as well as demographic, clinical, electrocardiographic, echocardiographic, and laboratory results.

The CHADS₂ score was the sum of 1 point each for heart failure (HF), HT, age ≥ 75 years, and DM and 2 points for prior stroke or transient ischemic attack. The CHA₂DS₂-VASC score was calculated by assigning 1 point each for CHF, HT, age 65–74 years, DM, vascular disease, and female gender and 2 points each for prior stroke or TIA and age ≥ 75 years. The newly formulated CHA₂DS₂-VASC-FSH score consists of CHF, HT, age ≥ 75 years (double score), DM, previous stroke/TIA (double score), vascular disease, age 65–74 years, sex category (male sex), hyperlipidaemia, smoking, and family history of CAD.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

The exclusion criteria included asymptomatic AF cases, cardiovascular arrest or asystole, any wide complex tachyarrhythmia presumed or suspected to be of ventricular origin, persistent (> 20 minutes) ST segment elevation; electrical cardioversion (pharmacological and spontaneous cardioversion were permissible) or chest compressions causing troponin I elevation; and recent history of myocardial infarction in the last 15 days. All study data were accessed via XXXXX University Hospital's automation system.

Statistical Analysis

Continuous study data were presented as mean \pm standard deviation or median (minimum-maximum) depending on the normality of their distribution assessed by Kolmogorov-Smirnov test. Categorical variables were presented as number and percentage. The difference between the groups with respect to continuous variables was tested with the Student's t test or Mann Whitney-U test, with the former being used for normally distributed variables and the latter for non-normally distributed ones. The difference with regard to categorical variables was tested using the Chi-squared test. Two separate univariate analyses were then performed to determine the correlation between examined study parameters and severe CAD and unstable CAD on coronary angiography. A binary logistic regression analysis was performed to determine the significant independent predictors of severe coronary artery disease on coronary angiography.

Results: The overall study population consisted of 72 patients with a mean age of 65.7 \pm 11.2 years, of whom 43 were male and 29 were female. Thirty-five (48.6%) patients had new-onset AF; 13 (18.1%) patients had paroxysmal AF; 4 (5.6%) had persistent AF; and 20 (27.8%) had permanent AF. Twenty-six (36.1%) patients had a history of coronary artery disease, with 6 (8.3%) having a history of stent implantation, 7 (9.7%) having a history of CABG operation, and 2 (2.8%) having a history of both stent implantation and CABG operation. Fifty-two (72.2%) patients presented with chest pain, of whom 46 (63.9%) had typical chest pain (**Table 1**).

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

A total of 30 (41.7%) patients had admission troponin I positivity; 44 (61.1%) had follow-up troponin I positivity, 50 (69.4%) patients had overall troponin I positivity, and 38 (52.8%) patients had a clinically significant delta troponin I change.

Thirty-seven (51.4%) patients were not found to have severe coronary lesions. A total of 35 (48.6%) patients had severe single- or multi-vessel coronary artery disease on coronary angiography. Twenty-four (68.6%) patients with severe coronary lesions underwent coronary intervention. A total of 24 (33.3%) patients underwent coronary intervention. Among these, 7 (29.2%) underwent coronary artery bypass grafting 17 (70.8%) underwent percutaneous coronary intervention and stenting (**Table 2**).

The comparison of the two groups with and without severe CAD revealed that the significant CAD group had significantly greater rates of hypertension, diabetes mellitus, CAD, CVA, severe renal disease, chest pain as the presenting symptom, and unstable lesions ($p < 0.05$ for all comparisons); significantly greater CHA₂DS₂-VASC ($p < 0.01$) and CHA₂DS₂-VASC-FSH scores ($p < 0.05$), and significantly greater RDW and creatinine levels ($p < 0.05$ for both). The two groups did not significantly differ with respect to cardiac biomarker levels, any troponin I positivity, ischemic ECG changes, echocardiographic wall motion abnormalities or left ventricular hypertrophy (**Table 3**).

On univariate analysis, the rate of severe CAD was significantly correlated to CHA₂DS₂VASC score ($p < 0.05$), CHA₂DS₂VASC-FSH score ($p < 0.05$), hypertension ($p < 0.05$), coronary artery disease (CAD) ($p < 0.05$), chronic severe renal disease ($p < 0.005$), chest pain as the presenting symptom ($p < 0.01$), creatinine ($p < 0.01$), and RDW ($p < 0.01$). Binary logistic regression analysis revealed that the CHA₂DS₂VASC-FSH score (OR=1.80 (95% CI 1.24-2.61; $p < 0.05$) and severe chronic renal disease (OR=7.96 (95% CI 1.46-43.51; $p < 0.05$) were the only significant predictors of severe coronary artery disease (**Table 4**). Among the three CAD prediction scores, the CHADS₂ score had an area under the curve (AUC) of 0.689 (0.567-0.811), the CHA₂DS₂VASC score had an AUC of 0.690 (0.568-0.813), and the CHA₂DS₂VASC-FSH score an AUC of 0.703

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

(0.582-0.825) A CHA₂DS₂VASc-FSH score of 3 or greater had a sensitivity of 77.1% and a specificity of 56.8% for predicting severe CAD (**Figure 1**).

The same correlation analyses were performed for patients with new-onset and paroxysmal AF (n=48). In univariate analysis, severe coronary lesions were correlated to CHA₂DS₂VASc score (p<0.05), CHA₂DS₂VASc-FSH score (p<0.05), left ventricular hypertrophy (p<0.05), history of coronary artery disease (p<0.05), chronic severe renal disease (p<0.05), HT (p<0.05), creatinine (p<0.05), and chest pain as the chief presenting symptom (p<0.05). On binary logistic regression analysis the CHA₂DS₂VASc-FSH score (OR=3.03 (95% CI 1.19-7.63); p<0.05) remained the only significant predictor of severe coronary artery disease (**Table 4**). Among the three scores, the CHADS₂ score had an AUC of 0.670 (0.514-0.827); CHA₂DS₂VASc score had an AUC of 0.671 (0.516-0.826), and the CHA₂DS₂VASc-FSH score had an AUC of 0.679 (0.517-0.842). A CHA₂DS₂VASc-FSH score of greater than 3 had a sensitivity of 80% and a specificity of 60.9% for predicting severe CAD (**Figure 2**).

Discussion

Atrial fibrillation (AF) may complicate an acute coronary syndrome [13,14] or it may also cause myocardial injury or infarction by creating myocardial supply-demand mismatch (type 2 myocardial ischemia), particularly when associated with rapid ventricular response, or by coronary embolism [15,16]. Unfortunately, AF may also be associated with electrocardiographic ST-T changes and troponin positivity mimicking myocardial ischemia and injury in the absence of severe CAD [2]. It is often difficult to discern AF-induced pseudo-infarction/ischemia patterns from type 2 myocardial ischemia resulting from underlying severe CAD, or a true acute coronary syndrome plus AF. Among patients with AF and unstable signs and symptoms, clinicians should determine the pretest probability of underlying severe or unstable CAD in order to make decisions about hospitalization and invasive coronary angiography.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaplar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

So far, only a few studies have studied the prevalence and predictors of severe CAD in patients with AF and unstable symptoms and found prevalences ranging from 18.2% to 32.5% [17,18]. Androulakis A et al [17] reported that there was no significant correlation between ischemic ECG changes and underlying occult CAD in patients with atrial fibrillation paroxysms. Alghamry et al [18] found that dyspnea, ST depression and history of coronary artery disease predicted significant CAD in patients hospitalized for AF but concluded that troponin elevation was common among these patients and was not a reliable predictor for underlying severe coronary lesions. Our study revealed that among patients with AF and unstable symptoms the prevalence of severe CAD was 48.6%, and it revealed that neither cardiac biomarkers nor ischemic electrocardiographic changes were able to predict underlying severe CAD. On the other hand, CHA₂DS₂VASc-FSH score and severe chronic renal disease emerged as significant independent predictors of severe CAD. Our results suggest that among patients presenting with AF and unstable symptoms, severe CAD can be independently predicted by the CHA₂DS₂VASc-FSH score irrespective of the type of AF, so that a score of 3 or greater had a relatively fair sensitivity and a moderate specificity for severe CAD. Whereas CHA₂DS₂-VASc score is originally designed to assess the risk of ischemic stroke and other thromboembolic events in AF [10], it gives an overall estimate of overall atherosclerotic burden on patients with AF since it incorporates many, albeit not all, risk factors for CAD. According to the Framingham study, AF and severe CAD share similar risk factors [19]. Given the multitude of common risk factors for both stroke and CAD, it has been suggested that CHA₂DS₂-VASc may provide prognostic information in CHD without known AF [20]. In support of this view, the CHA₂DS₂-VASc score has been used for determining CAD severity in various populations other than in AF setting. Cetin et al [7] reported that CHADS₂, CHA₂DS₂-VASc, and especially CHA₂DS₂-VASc-HS scores could be considered predictive of the risk of severe CAD determined by the Gensini score among patients undergoing diagnostic coronary angiography who were in sinus rhythm. In a similar study, Modi et al [9] reported that in 2976 consecutive patients, who were in sinus

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

rhythm and underwent coronary angiography, CHADS₂, CHA₂DS₂-VAsC, and especially CHA₂DS₂-VAsC-HS and CHA₂DS₂-VAsC-HSF scores were predictive of the risk of severe CAD. Hioki et al [20] reported that the CHA₂DS₂-VAsC score was able to predict MACE at 1 year in patients undergoing percutaneous coronary intervention who were in sinus rhythm. Subsequently, a number of previous studies used the modifications of the original CHA₂DS₂-VAsC for predicting the CAD severity in various settings. Çetin M et al [7] reported that the CHA₂DS₂-VAsC-HS score, which added hyperlipidemia and smoking to the original score, was significantly better at predicting severe CAD, assessed by the Gensini score and the number of diseased vessels, than the original CHADS₂ and CHA₂DS₂-VAsC scores in 407 consecutive patients undergoing coronary angiography. Uysal et al [21] used the CHA₂DS₂-VAsC-FSH score in patients with ST elevation myocardial infarction. The authors reported that the score was significantly correlated to atherosclerosis severity assessed by the SYNTAX score. Our study is noteworthy because it is the first to use the CHA₂DS₂-VAsC-FSH score in a population having AF and unstable signs and symptoms suggesting CAD. We used the newly developed CHA₂DS₂-VAsC-FSH score instead of the CHA₂DS₂-VAsC score because the former incorporates a greater number of risk factors for CAD and thus provides a greater clue about the underlying CAD than the traditional CHA₂DS₂-VAsC score. Supporting this idea, the CHA₂DS₂-VAsC score was correlated to severe CAD in univariate analysis but it did not independently predict severe CAD in multivariate analysis whereas the CHA₂DS₂-VAsC-FSH score did. The CHA₂DS₂-VAsC-FSH score was comparably predictive of significant coronary artery disease in early-stage AF types such as new-onset and paroxysmal AF, and the older AF types such as persistent and permanent AF. This suggests that it is the combination of risk factors and thus a greater CAD burden, but not the age of AF, that plays a role in development of CAD, and the CHA₂DS₂-VAsC-FSH score is equally predictive of severe underlying CAD in the overall AF population with unstable symptoms. It should also be noted that CHA₂DS₂-VAsC-FSH score helps provide the risk stratification not just in AF, but in all cases of ischemic

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VAsC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

heart disease, and it can be used in all of ischemic heart disease for prognostication. Therefore its role as a prognostic sign should be further studied.

This study has some limitations. First, it was a retrospective analysis and therefore is limited by the inherent limitations of its design. Hence, it involved only AF patients who underwent coronary angiography, and therefore the prevalence and the significant predictors of CAD remained unclear in the overall AF population including the conservatively managed ones. Furthermore, we lacked information about in-hospital or 30-day mortality and thus could not ascertain the prognostic value of CHA₂DS₂VASc-FSH score or delta troponin I change beyond coronary lesion characteristics. There is also a need for the validation of the CHA₂DS₂VASc-FSH in other AF settings, such as in outpatients, postoperative period, and ST elevation myocardial infarction.

In conclusion, in patients with AF and unstable symptoms warranting further management, the newly defined CHA₂DS₂VASc-FSH score can successfully predict severe CAD on coronary angiography. In clinical settings where clinical doubt exists whether symptoms are caused by severe CAD, or when severe CAD should be excluded for other purposes, a CHA₂DS₂VASc-FSH score of at least 3 can be used to make a decision to proceed with invasive coronary angiography. In the future the CHA₂DS₂VASc-FSH score can be incorporated with other biochemical and echocardiographic predictors of atherosclerosis in patients with AF.

Conflict of Interest: The authors declare no conflict of interest.

References:

1. Chenoweth J, Diercks DB. Management of atrial fibrillation in the acute setting. *Curr Opin Crit Care*. 2012;18:333-40.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçâğlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. *Eurasian J Med* 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

2. Tanindi A, Cemri M. Troponin elevation in conditions other than acute coronary syndromes. *Vasc Health Risk Manag.* 2011;7:597–603. (PMID:22102783).
3. Henriksson KM, Farahmand B, Johansson S, Asberg S, Terént A, Edvardsson N. Survival after stroke — the impact of CHADS2 score and atrial fibrillation. *Int J Cardiol*, 2010;141:18–23.
4. Biancari F, Asim Mahar MA, Kangasniemi OP. CHADS(2) and CHA(2)DS(2)-VASc scores for prediction of immediate and late stroke after coronary artery bypass graft surgery. *J Stroke Cerebrovasc Dis.* 2013;22:1304–1311.
5. Welles CC, Whooley MA, Na B, Ganz P, Schiller NB, Turakhia MP. The CHADS2 score predicts ischemic stroke in the absence of atrial fibrillation among subjects with coronary heart disease: data from the Heart and Soul Study. *Am Heart J.* 2011;162:555–561.
6. Poci D, Hartford M, Karlsson T, Herlitz J, Edvardsson N, Caidahl K. Role of the CHADS2 score in acute coronary syndromes: risk of subsequent death or stroke in patients with and without atrial fibrillation. *Chest.* 2012;141:1431–1440.
7. Cetin M, Cakici M, Zencir C, Tasolar H, Baysal E, Balli M, Aktürk E. Prediction of coronary artery disease severity using CHADS2 and CHA2DS2-VASc scores and a newly defined CHA2DS2-VASc-HS score. *Am J Cardiol.* 2014;113:950-6.
8. Lu DY, Huang CC, Huang PH, Chen JW, Chen TJ, Lin SJ, Chan WL, Lee CY, Leu HB. Usefulness of the CHADS2 Score for Prognostic Stratification in Patients With Coronary Artery Disease Having Coronary Artery Bypass Grafting. *Am J Cardiol.* 2017;119:839-844.
9. Modi R, Patted SV, Halkati PC, Porwal S, Ambar S, Mr P. Metgudmath V, Sattur A. et al. CHA2DS2-VASc HSF score New predictor of severity of coronary artery disease in 2976 patients. *Int J Cardiol* 2017;228:1002-1006.
10. January CT, Wann LS, Alpert JS, Calkins H, Cigarroa JE, Cleveland JC Jr, Conti JB, Ellinor PT, Ezekowitz MD, Field ME, et al. American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2014

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaylar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. *Eurasian J Med* 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol* 2014;64:e1-76.

11. Camm AJ, Kirchhof P, Lip GY, Schotten U, Savelieva I, Ernst S, Van Gelder IC, Al-Attar N, Hindricks G., Prendergast B, et al; ESC Committee for Practice Guidelines. Guidelines for the management of atrial fibrillation: the Task Force for the Management of Atrial Fibrillation of the European Society of Cardiology (ESC). European Heart Rhythm Association; European Association for Cardio-Thoracic Surgery. *Europace* 2010;12:1360-420.
12. Thygesen K, Mair J, Giannitsis E, Mueller C, Lindahl B, Blankenberg S, Huber K, Plebani M, Biasucci LM, Tubaro M, et al. How to use high-sensitivity cardiac troponins in acute cardiac care. *Eur Heart J* 2012;33:2252-2257.
13. Schmitt J. Atrial fibrillation in acute myocardial infarction: a systematic review of the incidence, clinical features and prognostic implications. *Eur Heart J* 2009;30:1038–1045.
14. Crenshaw BS, Ward SR, Granger CB, Stebbins AL, Topol EJ, Califf RM. Atrial fibrillation in the setting of acute myocardial infarction: the GUSTO-I experience. *Global Utilization of Streptokinase and TPA for Occluded Coronary Arteries. J Am Coll Cardiol* 1997;30:406–413.
15. Sandoval Y, Smith SW, Thorsen SE, Apple FS. Supply/demand type 2 myocardial infarction: should we be paying more attention? *J Am Coll Cardiol* 2014;63:2079–2087..
16. Kolodgie FD, Virmani R, Finn AV, Romero ME. Embolic myocardial infarction as a consequence of atrial fibrillation: a prevailing disease of the future. *Circulation* 2015;132:223–226.
17. Androulakis A, Aznaouridis KA, Aggeli CJ, Roussakis GN, Michaelides AP, Kartalis AN, Stougiannos PN, Dilaveris PE, Misovoulos PI, Stefanadis CI,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. *Eurasian J Med* 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

- et al. Transient ST-segment depression during paroxysms of atrial fibrillation in otherwise normal individuals: relation with underlying coronary artery disease. *J. Am. Coll. Cardiol* 2007;50:1909–1911.
18. Alghamry A, Hanna J, Pelecanos A, Kyranis S, Khelgi V, O'Rourke Patients, Carroll O, Oxenford C, Ranganwamaiah S, Tan C. Predictors of significant coronary artery disease in atrial fibrillation: Are cardiac troponins a useful measure. *Int J Cardiol* 2016;223: 744-749.
 19. Benjamin E.J, Levy D, Vaziri SM, D'Agostino RB, Belanger AJ, Wolf PA. Independent risk factors for atrial fibrillation in a population based cohort. The Framingham Heart Study. *JAMA* 1994;271:840–844.
 20. Hioki H, Miura T, Miyashita Y, Motoki H, Shimada K, Kobayashi Measurement, Nakajima H, Kimura H, Mawatari E, Akanuma H, et al. Risk stratification using the CHA₂DS₂-VASc score in patients with coronary heart disease undergoing percutaneous coronary intervention; sub-analysis of SHINANO registry. *IJC Heart & Vasculature* 2015;7:76-81.
 21. Uysal OK, Turkoglu C, Duran M, Kaya MG, Sahin DY, Gur M, Caylı M. Predictive value of newly defined CHA₂DS₂-VASc-HSF score for severity of coronary artery disease in ST segment elevation myocardial infarction. *Kardiol Pol* 2016;74:954-60.

Table 1. Demographic, Clinical, Angiographic characteristics of the study group

Characteristic	N(%)
Sex (male)	43 (59.7%)
Age (years)	65.7 ±11.2

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özın B, Müderrisoğlu İH. The Novel CHA₂DS₂-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. *Eurasian J Med* 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

Atrial fibrillation type	
New-onset	35 (48.6%)
Paroxysmal	13 (18.1%)
Persistent	4 (5.6%)
Permanent	20 (27.8%)
History of coronary artery disease	23 (31.9%)
PCI (stenting)	6 (8.3%)
CABG	7 (9.7%)
CABG+stenting	2 (2.8%)
Medically treated	8 (11.1%)
Diabetes mellitus	25 (34.7%)
Hypertension	48 (66.7%)
Smoking	26 (36.1%)
Hyperlipidemia	31 (43.1%)
Severe Renal disease	13 (18.1%)
Receiving Dialysis	6 (8.3%)
Not on dialysis	7(9.7%)
Heart failure	12 (16.7%)
Chronic obstructive pulmonary disease	9 (12.5%)
Thyroid disease	5 (6.9%)
History of cerebrovascular accident	7 (9.7%)
History of peripheral embolism	2 (2.8%)
Family history for coronary artery disease	10 (13.9%)
CHA2DS2-VASc score	3 (0-6)
CHA2DS2-VASc score \geq 2	55 (76.4%)
CHA2 DS2 -VASc-HSF score	4 (0-8)
Medications used	
Beta blocker	17(23.6%)
Non-dihydropyridine calcium channel blocker	7 (9.7%)
Digoxin	7 (9.7%)
Propafenone	2 (2.8%)
Amiodarone	1 (1.4%)
Warfarin Na	7 (9.7%)
NOAC	3 (4.2%)
Aspirin	21 (29.2%)
Clopidogrel	3 (4.2%)
ACE inhibitor	10 (13.9%)
ARB	11 (15.3%)
Dihydroopyridine calcium channel blocker	6 (8.4%)
Diuretic	19 (26.4%)
Insulin	6 (8.4%)
Oral antidiabetics	14 (19.4%)
Statins	6 (8.4%)
Thyroid medication	2 (2.8%)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

Table-2 Findings on coronary angiography and coronary intervention results

Coronary angiography findings	
Severe coronary artery disease	35 (48.6%)
Single vessel disease	8 (22.9%)
Multivessel disease	27 (77.1%)
Non-severe coronary artery disease	37 (51.4%)
Normal coronary arteries	14 (37.8%)
Luminal irregularities	13 (35.1%)
Non-significant single vessel	3 (8.1%)
Non-significant multivessel	2 (5.4%)
Patent stent(s)	1 (2.7%)
Patent bypass graft(s)	4 (5.6%)
Unstable lesions	11 (15.3%)
Thrombosed lesion(s)	8 (22.9%)
Dissection	2 (5.4%)
Markedly slow flow	1 (2.7%)
Coronary intervention	24 (33.3%)
PCI	17 (70.8%)
CABG	7 (9.7%)
Stented vessel (or major branch(es) of)	
LMCA	1 (5.9%)
LAD	4 (23.5%)
Cx	6 (35.3%)
RCA	5 (29.4%)
2-vessel coronary disease	1 (5.9%)
Bypassed vessel (or major branch(es) of)	
LAD	6 (8.4%)
Cx	4 (5.6%)
RCA	4 (5.6%)

PCI: percutaneous coronary intervention, CABG: coronary artery bypass grafting, LMCA: Left main coronary artery, LAD: Left anterior descending artery, Cx: Circumflex artery, RCA: right coronary artery

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

Table 3. Comparison of the study groups with respect to demographic, biochemical, electrocardiographic, and echocardiographic variables

Parameter	Severe CAD (+) (n=35)	Severe CAD (-) (n=37)	p
Sex (male)	21 (60%)	22 (59.5%)	0.963
Age (years)	68.34 (11.74)	63.3 (10.3)	0.058
AF type			
New-onset	21 (60%)	14 (37.8%)	0.112
Paroxysmal	4 (11.4%)	9 (24.3%)	
Persistent	0 (0%)	4 (10.8%)	
Permanent	10 (28.6%)	10 (27.0%)	
CAD	17 (48.6%)	7 (18.9%)	<0.05
DM	18 (51.4%)	7 (18.9%)	<0.05
HT	29 (82.9%)	19 (51.3%)	<0.05
Smoking	12 (34.3%)	14 (37.1%)	0.627
HL	19 (54.3%)	12 (32.4%)	0.063
CRD	12 (34.3%)	1 (2.7%)	<0.05
Heart failure	8 (22.9%)	4 (10.8%)	0.467
COPD	3 (8.6%)	7 (18.9%)	0.458
Thyroid disease	3 (8.6%)	2 (5.4%)	0.584
CVA	6 (17.1%)	1 (2.7%)	<0.05
History of peripheral embolism	1 (2.9%)	0 (0%)	0.782
CHA2DS2-VASc score	4 (0-6)	2 (0-5)	<0.01
CHA2 DS2 -VASc-HSF score	5 (1-8)	3 (0-7)	<0.05
Family history for CAD	4 (11.4%)	6 (16.2%)	0.563
Chest pain as the chief symptom	30 (85.7%)	22 (59.5%)	<0.05
Typical angina pectoris	26 (74.3%)	20 (54.1%)	0.075
Unstable lesions	11 (31.4%)	0 (0%)	<0.01
Admission troponin I (ng/ml)	0.91±3.08	1.57 ±4.64	0.491
Follow-up Troponin I (ng/ml)	0.42 (0-370.80)	0.06 (0-119.31)	0.428
Admission troponin I positivity	16 (45.7%)	14 (37.8%)	0.567
Follow-up troponin I positivity	25 (71.4%)	19 (51.4%)	0.166
Overall troponin I positivity	22 (62.9%)	20 (54.1%)	0.099
Delta troponin I change	23 (65.7%)	15 (40.5%)	0.078
Admission CK/MB (ng/ml)	3.62 ±3.65	3.33 ±4.17	0.759
Follow-up CK/MB (ng/ml)	7.59 ±4.55	5.33 ±11.49	0.150
Blood Urea Nitrogen (mg/dl)	20.66 ±7.82	21.14 ± 12.94	0.855

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçağlar E, Yılmaz M, Özın B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

Creatinine (mg/dl)	1.06 (0.61-5.31)	0.90 (0.66-4.45)	<0.05
Sodium (mEq/L)	137.34 ±4.11	137.78 ±3.19	0.631
Potassium (mEq/L)	3.86 ±1.31	4.05 ±0.85	0.487
Hemoglobin count (g/dl)	12.90 ±3.86	13.28 ±3.79	0.684
Leucocyte count (10 ³ /μl)	8.31 ±3.36	9.16 ±3.82	0.641
Neutrophil count (10 ³ /μl)	5.98 (2.06-38.21)	5.32 (3.25-19.50)	0.741
Platelet count (10 ³ /μl)	222.54 ±92.22	222.34 ±73.12	0.992
Lymphocyte count (10 ³ /μl)	2.10 ±1.16	2.33 ±2.62	0.232
NLR	3.59 ±2.76	3.83 ±3.14	0.098
PLR	130.43 ±73.86	131.18 ±72.59	0.967
RDW (%)	15.7 (12.0-21.7)	14.5(11.5-22.7)	<0.05
Ischemic changes on ECG	14 (40.0%)	8 (21.6%)	0.095
Heart rate on ECG (bpm)	106.8±29.7	111.8±33.2	0.088
WMA	19 (54.3%)	13 (35.1%)	0.232
LVEF (%)	46.3 ±11.8)	49.8 ±11.9)	0.229
LVHT	25 (71.4%)	20 (54.1%)	0.192

CAD: Coronary artery disease, DM: diabetes mellitus, HT: Hypertension, HL: hyperlipidemi, CRD: Chronic renal disease, COPD: chronic obstructive pulmonary disease, CVA: cerebrovascular accident, NLR: neutrophil-to-lymphocyte ratio, PLR: platelet-lymphocyte ratio, RDW: red cell distribution width, ECG: electrocardiogram, LVEF: left ventricular ejection fraction, LVHT: left ventricular hypertrophy, WMA: Wall motion abnormality on echocardiography

Table 4. Predictors of severe CAD on coronary angiography in the overall AF and new-onset AF populations

		Univariate analysis (p)	Binary logistic regression analysis with Backward Wald method (OR (95% CI)
Overall AF	Severe	CHA2DS2VASC-FSH (<0.05)	CHA2DS2VASC-FSH: 1.80 (1.24-2.61) CRD: 7.96 (1.46-43.51)
	CAD	CHA2DS2VASC (<0.05) HT (<0.05) CAD (<0.05) CRD (<0.05) Chest pain (<0.01) Creatinine (0 <0.01) RDW (<0.01)	
		CHA2DS2VASC-FSH (<0.05) CHA2DS2VASC (0<0.05) LVHT (<0.05), CAD (<0.05)	CHA2DS2VASC-FSH: 3.03 (1.19-7.63)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaglar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org

New-onset AF	Severe CAD	CRD (<0.05) HT (<0.05) Creatinine (<0.05) Chest pain (<0.05)	
--------------	------------	---	--

AF: atrial fibrillation, CAD: coronary artery disease, HT: hypertension, CRD: chronic renal disease

Figure Legends:

Figure 1. ROC curves of the thromboembolic risk scores in the prediction of severe coronary artery disease in the overall atrial fibrillation group.

Figure 2. ROC curves of the thromboembolic risk scores in the prediction of severe coronary artery disease in the combined new-onset and paroxysmal atrial fibrillation groups.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as: Çiftçi O, Yılmaz KC, Karaçaplar E, Yılmaz M, Özin B, Müderrisoğlu İH. The Novel CHA2DS2-VASC-FSH Score is Predictive of Severe Coronary Artery Disease on Coronary Angiography in Patients with Atrial Fibrillation and Unstable Symptoms. Eurasian J Med 2019; DOI: 10.5152/eurasianjmed.2019.18342. ©Copyright 2019 by Atatürk University School of Medicine - Available online at www.eajm.org